

Local Coverage Determination (LCD): MoIDX: MGMT Promoter Methylation Analysis (L36192)

Links in PDF documents are not guaranteed to work. To follow a web link, please use the MCD Website.

Contractor Information

Contractor Name	Contract Type	Contract Number	Jurisdiction	State(s)
Noridian Healthcare Solutions, LLC	A and B MAC	02101 - MAC A	J - F	Alaska
Noridian Healthcare Solutions, LLC	A and B MAC	02102 - MAC B	J - F	Alaska
Noridian Healthcare Solutions, LLC	A and B MAC	02201 - MAC A	J - F	Idaho
Noridian Healthcare Solutions, LLC	A and B MAC	02202 - MAC B	J - F	Idaho
Noridian Healthcare Solutions, LLC	A and B MAC	02301 - MAC A	J - F	Oregon
Noridian Healthcare Solutions, LLC	A and B MAC	02302 - MAC B	J - F	Oregon
Noridian Healthcare Solutions, LLC	A and B MAC	02401 - MAC A	J - F	Washington
Noridian Healthcare Solutions, LLC	A and B MAC	02402 - MAC B	J - F	Washington
Noridian Healthcare Solutions, LLC	A and B MAC	03101 - MAC A	J - F	Arizona
Noridian Healthcare Solutions, LLC	A and B MAC	03102 - MAC B	J - F	Arizona
Noridian Healthcare Solutions, LLC	A and B MAC	03201 - MAC A	J - F	Montana
Noridian Healthcare Solutions, LLC	A and B MAC	03202 - MAC B	J - F	Montana
Noridian Healthcare Solutions, LLC	A and B MAC	03301 - MAC A	J - F	North Dakota
Noridian Healthcare Solutions, LLC	A and B MAC	03302 - MAC B	J - F	North Dakota
Noridian Healthcare Solutions, LLC	A and B MAC	03401 - MAC A	J - F	South Dakota
Noridian Healthcare Solutions, LLC	A and B MAC	03402 - MAC B	J - F	South Dakota
Noridian Healthcare Solutions, LLC	A and B MAC	03501 - MAC A	J - F	Utah
Noridian Healthcare Solutions, LLC	A and B MAC	03502 - MAC B	J - F	Utah
Noridian Healthcare Solutions, LLC	A and B MAC	03601 - MAC A	J - F	Wyoming
Noridian Healthcare Solutions, LLC	A and B MAC	03602 - MAC B	J - F	Wyoming

[Back to Top](#)

LCD Information

Document Information

LCD ID L36192	Original Effective Date For services performed on or after 04/15/2016
LCD Title MoIDX: MGMT Promoter Methylation Analysis	Revision Effective Date For services performed on or after 01/01/2018
Proposed LCD in Comment Period N/A	Revision Ending Date N/A
Source Proposed LCD N/A	Retirement Date N/A
AMA CPT / ADA CDT / AHA NUBC Copyright Statement	Notice Period Start Date 02/25/2016
	Notice Period End Date 04/14/2016

CPT only copyright 2002-2018 American Medical Association. All Rights Reserved. CPT is a registered trademark of the American Medical Association. Applicable FARS/DFARS Apply to Government Use. Fee schedules, relative value units, conversion factors and/or related components are not assigned by the AMA, are not part of CPT, and the AMA is not recommending their use. The AMA does not directly or indirectly practice medicine or dispense medical services. The AMA assumes no liability for data contained or not contained herein.

The Code on Dental Procedures and Nomenclature (Code) is published in Current Dental Terminology (CDT). Copyright © American Dental Association. All rights reserved. CDT and CDT-2016 are trademarks of the American Dental Association.

UB-04 Manual. OFFICIAL UB-04 DATA SPECIFICATIONS MANUAL, 2014, is copyrighted by American Hospital Association ("AHA"), Chicago, Illinois. No portion of OFFICIAL UB-04 MANUAL may be reproduced, sorted in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without prior express, written consent of AHA." Health Forum reserves the right to change the copyright notice from time to time upon written notice to Company.

CMS National Coverage Policy

Title XVIII of the Social Security Act (SSA), §1862(a)(1)(A), states that no Medicare payment shall be made for items or services that "are not reasonable and necessary for the diagnosis or treatment of illness or injury or to improve the functioning of a malformed body member."

Title XVIII of the Social Security Act, §1833(e), prohibits Medicare payment for any claim lacking the necessary documentation to process the claim.

42 Code of Federal Regulations (CFR) §410.32 Diagnostic x-ray tests, diagnostic laboratory tests, and other diagnostic tests: Conditions.

CMS Internet Online Manual Pub. 100-02 (Medicare Benefit Policy Manual), Chapter 15, Section 80, "Requirements for Diagnostic X-Ray, Diagnostic Laboratory, and Other Diagnostic Tests".

CMS Internet-Only Manuals, Publication 100-04, Medicare Claims Processing Manual, Chapter 16, §50.5 Jurisdiction of Laboratory Claims, 60.12 Independent Laboratory Specimen Drawing, 60.2. Travel Allowance.

CMS Internet Online Manual Pub. 100-04 (Medicare Claims Processing Manual), Chapter 23 (Section 10) "Reporting ICD Diagnosis and Procedure Codes".

CMS Internet-Only Manual, Pub 100-04, Medicare Claims Processing Manual, Chapter 12, §30-Correct Coding Policy.

Coverage Guidance

Coverage Indications, Limitations, and/or Medical Necessity

Indications and Limitations of Coverage

This policy provides limited coverage for methylation analysis for hypermethylation of the O-6--methylguanine DNA methyltransferase (MGMT) gene promoter. MGMT methylation analysis testing is considered to be reasonable and necessary for adult patients when the following criteria are met:

- Tumor type is high--grade malignant glioma (e.g. glioblastoma multiforme (GBM), anaplastic astrocytoma) **and**

- Patients are able to tolerate temozolomide therapy or radiation therapy, **and**
- The physician will use the MGMT testing results to decide between radiation therapy and chemotherapy alone as 1st line adjuvant treatment, or between temozolomide and other chemotherapy for 1st line adjuvant treatment

Note: This assessment is predicated on the assumption that therapy is considered beneficial for the specific patient.

Summary of Evidence

Cancer is the consequence of genetic alterations that result in a deregulation of important cellular pathways responsible for various essential functions, including cell growth, cell cycle progression, and apoptosis (programmed cell death). One result of these genetic alterations is gliomas. The treatment of high-grade gliomas, especially GBM, remains difficult as no contemporary treatments are curative. For the past several years, the standard treatment for GBM consists of maximal surgical resection, radiotherapy (RT), and concomitant and adjuvant chemotherapy with temozolomide.

Although surgical resection, RT, and chemotherapy with temozolomide are considered standard of care for most patients with high-grade glioma (including GBM and anaplastic astrocytoma), not all patients tolerate these treatments. For patients older than 70 years with a low performance rating, radiation or temozolomide alone is sometimes employed. Temozolomide treatment is not considered inferior to radiation therapy and may be tolerated better than RT by "frail" patients with low performances scores.

In patients for whom temozolomide is not the current standard of care, it has been proposed that MGMT methylation analysis can be used to predict the efficacy of temozolomide treatment. Epigenetic silencing of the MGMT (O-6-methylguanine-DNA methyltransferase) DNA repair gene, by promoter methylation, leads to a lack of MGMT protein expression. Lack of MGMT protein expression immunohistochemically is related to drug responses in patients with malignant glioma treated with alkylating agents. In particular, MGMT hypermethylation is a known predictive biomarker of response to temozolomide treatment with favorable outcomes in terms of overall survival (OS) and progression free survival (PFS) in GBM patients.

MGMT promoter methylation status is a strong and independent prognostic factor in patients with newly diagnosed GBM and a clinically relevant predictive marker in the subpopulation of elderly GBM patients. MGMT promoter methylation analysis can aid in treatment decisions for patients over 70. For patients older than 70 with a good performance rating, there is evidence of benefit of temozolomide in addition to RT. In patients with lower performance, temozolomide can be used alone as it was found to be equally as effective as RT alone and it has lower toxicity for the frail population. In the temozolomide arm of both the Nordic and German trials, patients with MGMT promoter methylation had longer survival than those without. (9.7 vs 6.8 months; HR, 0.56; 95% CI, 0.34-0.93)

MGMT promoter methylation analysis also has prognostic utility. However performing MGMT analysis is only recommended by NCCN guidelines for temozolomide guidance and not for overall prognosis prediction. Lattanzio et al confirmed that patients carrying methylation of the MGMT promoter reported a longer OS and PFS than patients with an unmethylated promoter. Wang et al also evaluated the prognostic value of MGMT promoter methylation and TP53 mutation status found similar results.

There is still a lack of consensus on the optimal assay for reliable MGMT promoter methylation testing and a variety of tests are being used in different laboratories. According to Berghoff et al, pyrosequencing is the only method for which an adequately high analytical performance (high intra- and inter-laboratory repeatability and reproducibility) has been demonstrated in a fully published trial. MGMT promoter methylation testing should be performed by an experienced laboratory in which this testing has been validated.

MGMT may also be useful for determining the prognosis of colorectal cancer patients and to identify those requiring more aggressive adjuvant therapies. Future studies will be necessary to determine its clinical utility in this area. Likewise, MGMT methylation may be an important biomarker in subsets of esophageal cancers where temozolomide may be utilized to successfully treat these patients, but where additional research on clinical utility is also needed. MGMT methylation analysis is also mentioned in the literature as a predictive marker for ovarian cancer and melanoma. However, evidence on the use of MGMT testing is unclear in these diagnoses and additional studies are needed on the clinical utility in these cancers.

Analysis of Evidence

(Rationale for Determination)

Level of Evidence:

Quality - Strong

Strenght - Strong

Weight - Moderate

In summary, the current literature and NCCN guidelines support the use of MGMT methylation analysis to predict the usefulness of temozolomide treatment in adult patients with high-grade gliomas.

[Back to Top](#)

Coding Information

Bill Type Codes:

Contractors may specify Bill Types to help providers identify those Bill Types typically used to report this service. Absence of a Bill Type does not guarantee that the policy does not apply to that Bill Type. Complete absence of all Bill Types indicates that coverage is not influenced by Bill Type and the policy should be assumed to apply equally to all claims.

022x Skilled Nursing - Inpatient (Medicare Part B only)

Revenue Codes:

Contractors may specify Revenue Codes to help providers identify those Revenue Codes typically used to report this service. In most instances Revenue Codes are purely advisory. Unless specified in the policy, services reported under other Revenue Codes are equally subject to this coverage determination. Complete absence of all Revenue Codes indicates that coverage is not influenced by Revenue Code and the policy should be assumed to apply equally to all Revenue Codes.

N/A

CPT/HCPCS Codes

Group 1 Paragraph: N/A

Group 1 Codes:

81287 MGMT (O-6-METHYLGUANINE-DNA METHYLTRANSFERASE) (EG, GLIOBLASTOMA MULTIFORME), METHYLATION ANALYSIS

ICD-10 Codes that Support Medical Necessity

Group 1 Paragraph: N/A

Group 1 Codes:

ICD-10 Codes

Description

C71.0	Malignant neoplasm of cerebrum, except lobes and ventricles
C71.1	Malignant neoplasm of frontal lobe
C71.2	Malignant neoplasm of temporal lobe
C71.3	Malignant neoplasm of parietal lobe
C71.4	Malignant neoplasm of occipital lobe
C71.5	Malignant neoplasm of cerebral ventricle
C71.6	Malignant neoplasm of cerebellum
C71.7	Malignant neoplasm of brain stem
C71.8	Malignant neoplasm of overlapping sites of brain

ICD-10 Codes	Description
C71.9	Malignant neoplasm of brain, unspecified

ICD-10 Codes that DO NOT Support Medical Necessity N/A

ICD-10 Additional Information [Back to Top](#)

General Information

Associated Information

Documentation Requirements

The patient's medical record must contain documentation that fully supports the medical necessity for services included within this LCD. (See "Coverage Indications, Limitations, and/or Medical Necessity") This documentation includes, but is not limited to, relevant medical history, physical examination, and results of pertinent diagnostic tests or procedures.

Documentation supporting the medical necessity should be legible, maintained in the patient's medical record, and must be made available to the MAC upon request.

Sources of Information

N/A

Bibliography

- Berghoff AS, Hainfellner JA, Marosi C, et al. Assessing MGMT methylation status and its current impact on treatment in glioblastoma. *CNS Oncol.* 2015 Jan;4(1):47-52.
- Chang IW, Hsu CT, Lin JW, et al. The prognostic impact of MGMT expression on low-grade gangliogliomas: a clinicopathological and immunohistochemical study. *Folia Neuropathol.* 2013;51(4):275-82.
- Furnari FB, Fenton T, Bachoo RM, et al. Malignant astrocytic glioma: genetics, biology, and paths to treatment. *Genes Dev.* Nov 1, 2007;21(21):2683-710.
- Hasina R1, Surati M, Kawada I, et al. O-6-methylguanine-deoxyribonucleic acid methyltransferase methylation enhances response to temozolomide treatment in esophageal cancer. *J Carcinog.* 2013 Oct 28;12:20. doi: 10.4103/1477-3163.120632. eCollection 2013.
- Hegi et al. MGMT Gene Silencing and Benefit from Temozolomide in Glioblastoma. 2005 *N Engl J Med* 352: 997-1003.
- Keime-Guibert F, Chinot O, Taillandier L, et al. Radiotherapy for glioblastoma in the elderly. *N Engl J Med.* Apr 12, 2007; 356(15): 1527-35.
- Krex D et al. German Glioma Network. Long-term survival with glioblastoma multiforme. *Brain.* 2007, 130:2596-606.
- Larijani L, Madjd Z, Samadikuchaksaraei A, et al. Methylation of O6-methyl guanine methyltransferase gene promoter in meningiomas comparison between tumor grades I, II, and III. *Asian Pac J Cancer Prev.* 2014;15(1):33-8.
- Lattanzio L, Borgognone M, Mocellini C, et al. MGMT promoter methylation and glioblastoma: a comparison of analytical methods and of tumor specimens. *Int J Biol Markers.* 2014 Dec 25:0.
- Malmstrom A, Gronberg BH, Marosi C, et al. Temozolomide versus standard 6-week radiotherapy versus hypofractionated radiotherapy in patients older than 60 years with glioblastoma: the Nordic randomised phase 3 trial. *Lancet Oncol* 2012. Sep;13(9):916-26.
- Martinez R et al. Frequent hypermethylation of the DNA repair gene MGMT in long-term survivors of glioblastoma multiforme. *J Neurooncol.* 2007 83(1):91-3

12. Mollemann M, Wolter M, Felsberg J, et al. Frequent promoter hypermethylation and low expression of the MGMT gene in oligodendroglial tumors. *Int J Cancer* 2005; 113:379-385.
13. NCCN Guidelines: Central Nervous System Cancers, Version 1.2015
14. Ogino S et al. Precision and performance characteristics of bisulfite conversion and real-time PCR (MethyLight) for quantitative DNA methylation analysis. *J Mol Diagn.* 2006;8(2):209-17.
15. Oliver JA, Ortiz R, Melguizo C1, et al. Prognostic impact of MGMT promoter methylation and MGMT and CD133 expression in colorectal adenocarcinoma. *BMC Cancer.* 2014 Jul 11;14:511. doi: 10.1186/1471-2407-14-511.
16. Preusser M, de Ribaupierre S, Wohrer A, et al. Current concepts and management of glioblastoma. *Ann Neurol.* Jul 2011;70(1):9-21.
17. Rizzo D, Scalzone M, Ruggiero A, et al. Temozolomide in the treatment of newly diagnosed diffuse brainstem glioma in children: a broken promise? *J Chemother.* 2014 Feb;27(2):106-10.
18. Roa W, Brasher PM, Bauman G, et al. Abbreviated course of radiation therapy in older patients with glioblastoma multiforme: a prospective randomized clinical trial. *J Clin Oncol.* May 1 2004;22(9):1583-8.
19. Sathornsumetee S, Reardon DA, Desjardins A, et al. Molecularly targeted therapy for malignant glioma. *Cancer.* Jul 1 2007;110(1):13-24.
20. Wang K, Wang YY, Ma J, et al. Prognostic Value of MGMT Promoter Methylation and TP53 Mutation in Glioblastomas Depends on IDH1 Mutation. *Asian Pac J Cancer Prev.* 2014;15(24):10893-8.
21. Zhang W, Lin Y, Chen B, et al. Recurrent glioblastoma of childhood treated with bevacizumab: case report and molecular features. *Childs Nerv Syst.* 2010 Jan;26(1):137-43.

[Back to Top](#)

Revision History Information

Revision History Date	Revision History Number	Revision History Explanation	Reason(s) for Change
01/01/2018	R1	This policy is revised to comply with the 21st Century Cures Act.	<ul style="list-style-type: none"> • Creation of Uniform LCDs With Other MAC Jurisdiction

[Back to Top](#)

Associated Documents

Attachments N/A

Related Local Coverage Documents Article(s) [A54896 - Response to Comments: MoIDX: MGMT Promoter Methylation Analysis](#) LCD(s) [DL36190](#) - (MCD Archive Site) [DL36192](#) - (MCD Archive Site)

Related National Coverage Documents N/A

Public Version(s) Updated on 12/28/2017 with effective dates 01/01/2018 - N/A [Updated on 02/05/2016 with effective dates 04/15/2016 - N/A](#) [Back to Top](#)

Keywords

- MoIDX
- MGMT
- Methylation
- methyltransferase

- methylguanine
- glioma
- glioblastoma
- GBM
- anaplastic
- astrocytoma
- temozolomide
- 81287

Read the [LCD Disclaimer](#) [Back to Top](#)